arXiv:1211.2580v1 [physics.soc-ph] 12 Nov 2012

# Quantifying Entanglement in Quantum Complex Networks

Alessio Cardillo,[1, 2] Fernando Galve,[3] David Zueco,[2, 4, 5] and Jesús Gómez-Gardeñes[1, 2]
[1] *Instituto de Biocomputación y Física de Sistemas Complejos,*
*Universidad de Zaragoza, E-50018 Zaragoza, Spain.*
[2] *Departamento de Física de la Materia Condensada,*
*Universidad de Zaragoza, E-50009 Zaragoza, Spain.*
[3] *Institute for Cross Disciplinary Physics and Complex Systems, IFISC (CSIC-UIB), Palma de Mallorca, Spain*
[4] *Instituto de Ciencia de Materiales de Aragón (ICMA),*
*CSIC-Universidad de Zaragoza, E-50012 Zaragoza, Spain.*
[5] *Fundación ARAID, Paseo María Agustín 36, E-50004 Zaragoza, Spain.*
(Dated: November 13, 2012)

We introduce the use of entanglement entropy as a tool for studying the amount of information stored in quantum complex networks. By considering the ground state of a network of coupled quantum harmonic oscillators, we compute the information that each node has on the rest of the system. We show both analytically and numerically that the nodes storing the largest amount of information are not the ones with the highest connectivity, but those with intermediate connectivity thus breaking down the usual hierarchical picture of classical networks. As a byproduct, our results point out that the amount of information available for an external controller connecting to a quantum network is bounded although it can be maximized by means of a moderate number of connections.

PACS numbers: 89.75.Fb,03.67.-a,89.70.Cf

The advent of network science has influenced the research in many fields of science in general, and physics in particular, in a pervasive way [1]. Since the discovery of the structural features of real social, biological and technological networks [2, 3], the development of the theoretical machinery of network science has blossomed as an efficient framework to interpret the many interaction patterns encoded in real-scale complex systems of diverse nature [4] and to model correctly the dynamical processes taking place on top of them [5, 6].

One of the most important avenues of research in network science is its connection with information theory. In this way, different information-theoretical tools have been proposed to characterize the complexity of networks beyond the typical statistical indicators such as their degree distribution, clustering coefficient, degree correlations, etc [4]. For instance, Shannon entropy, as shown in [7–10], has been successfully applied to characterize the complexity of ensembles of networks sharing some structural features while information-theoretical tools have been also applied to the study of diffusion processes on top of networks, such as random walks [11–16].

The synergy between the field of complex networks and that of information theory has recently appealed to the quantum information community. As a product, classical results on percolation theory [17–20] and network science, such as the small-world effect [21], have been revisited in networked structures of coupled quantum systems as a first step for designing quantum communication networks. Conversely, the use of quantum dynamical processes, such as quantum random walks [22] and their application to rank the importance of network elements [23–26], has given new quantum information perspectives to classical problems of the network realm.

In this Letter we quantify the amount of information that a single element of a quantum network shares with the rest of the system. To this aim we consider a network of quantum harmonic oscillators and analyze its ground state to compute the entropy of entanglement that vacuum fluctuations creates between single nodes and the rest of the network. By using the Von Neumann entropy [27, 28] we show that the amount of information available to nodes is limited, regardless of how large the number of connections of a node is. Moreover, in contrast to the usual scenario in classical networks, the nodes with the largest entanglement are not the hubs, but those components with intermediate connectivity. As a byproduct, our results show that in order to extract the maximum information of a quantum network it is enough to add an external controller connected to a moderate number of elements. Alternatively, a perturbation via quantum operations of intermediate-degree nodes induces the maximum damage to the quantum network state. Finally we round off the Letter by commenting on further implications of our results in quantum gravity models.

*Theory and Model.-* As usual, we define a network as a set of $N$ nodes and $E$ edges (or links) accounting for their pairwise interactions. The network backbone is usually encoded in the Adjacency matrix, $A$, such that $A_{ij} = 1$ if an edge connects nodes $i$ and $j$ while $A_{ij} = 0$ otherwise. In this Letter we restrict to undirected networks so that $A_{ij} = A_{ji}$. Although matrix $A$ stores all the structural meaning of a network it is more convenient to rely on the so-called network Laplacian, $L$, to analyze its structural and dynamical properties [29]. The Laplacian of a network is defined from the Adjacency matrix as $L_{ij} = k_i\delta_{ij} - A_{ij}$, where $k_i = \sum_j A_{ij}$ is the connectivity of node $i$, *i.e.*, the number of nodes connected to $i$.

Once defined the physical skeleton of the network we define its dynamical setup. As anticipated above, we consider nodes as identical quantum oscillators, interacting as dictated by the network topology encoded in $L$. The Hamiltonian of the quantum network thus reads:

$$H_{\rm network} = \frac{1}{2}\Big(\mathbf{p}^{\rm T}\mathbf{p} + \mathbf{x}^{\rm T}(\mathbb{I}+2cL)\mathbf{x}\Big)\,, \tag{1}$$

here $\mathbb{I}$ is the $N \times N$ identity matrix, $c$ is the coupling strength between connected oscillators while $\mathbf{p}^{\rm T} = (p_1, p_2, ..., p_N)$ and $\mathbf{x}^{\rm T} = (x_1, , x_2, ..., x_N)$ are the operators corresponding to the momenta and positions of nodes respectively, satisfying the usual commutation relations: $[\mathbf{x}, \mathbf{p}^{\rm T}] = i\hbar\mathbb{I}$ (we set $\hbar = 1$ in the following). The characterization of the quantum dynamical states is given by the eigenvectors of the Laplacian matrix and their corresponding eigenvalues shifted by 1. Note that, although the classical ground state of the above system is trivial (all the particles having $x_i = p_i = 0$), the quantum nature of the system prevents nodes from being at rest due to Heisenberg uncertainty principle. Moreover, the nature of the associated quantum vacuum fluctuations only depends on the pattern of physical interactions, *i.e.*, the network structure.

We now study the properties of the ground state to quantify the amount of information each of the elements of a network shares with the rest of the system via quantum fluctuations. To this aim, we consider the partition of the network into a node, say $i$, and its complement $i^{\rm c}$, *i.e.* the rest of the network. Then, we compute the mutual information shared by the two parties as:

$$\mathcal{I}(i|i^{\rm c}) = S_i + S_{i^{\rm c}} - S_{\rm tot}\,. \tag{2}$$

Here $S_i$ and $S_{i^{\rm c}}$ are marginal entropies and $S_{\rm tot}$ is the total entropy of the network. It is natural to choose the Von Neumann entropy to quantify the quantum information of the system, yielding $S_{\rm tot} = 0$ for the ground state (as it is a *pure state*). The marginal entropies for $i$ and $i^c$ read [30]:

$$S_i = S_{i^{\rm c}} = \Big(\mu_i + \frac{1}{2}\Big)\log\Big(\mu_i + \frac{1}{2}\Big) - \Big(\mu_i - \frac{1}{2}\Big)\log\Big(\mu_i - \frac{1}{2}\Big)\,, \tag{3}$$

which is a monotonically increasing function of $\mu_i$ that is characterized by the second moments of the positions and momenta of nodes, $\mu_i = \sqrt{\langle x_i^2\rangle\langle p_i^2\rangle}$. After some algebra (see Appendix A) we are able to quantify the value of $\mu_i$ as:

$$\mu_i^2 = \frac{1}{4}\sum_{j,j'} S_{ij}^2 S_{ij'}^2 \sqrt{\frac{1+2c\lambda_j}{1+2c\lambda_{j'}}}\,, \tag{4}$$

where $\{\lambda_j\}$ are the eigenvalues of the network Laplacian $L$ and matrix $S$ accounts for the normal mode transformation that diagonalizes the network Laplacian: $L_d = S^T L S$ with $S^T S = \mathbb{I}$.

From Eq.(3) it is clear that each node has some mutual information with the rest of the system provided $\mu_i > 1/2$ whereas from Eq.(4) we conclude that the amount of information depends on its contribution to each of the Laplacian eigenvectors. We finish by noting that, since the total network is in its ground (and pure) state we have $S_i = S_{i^{\rm c}} = \mathcal{I}(i|i^{\rm c})/2$. Therefore, the information that a node shares with the network is intrinsically due to quantum correlations. Equivalently, the mutual information is, itself, a measure of the entanglement (quantified by $S_i$) between a single node and the rest of the system.



FIG. 1. (color online) Average entropy $\langle S\rangle$ *versus* the coupling strength, $c$, for Random Regular Graphs (RRG) ●, Erdős-Rényi (ER) ■ networks, and Configurational (SF-CONF) ♦ and Barabási-Albert Scale-Free (SF-BA) ▲ networks. The average connectivity of the networks is $\langle k\rangle = 4$.

In the following we quantify the entanglement entropies of nodes embedded in different network topologies. First, we explore two homogeneous network substrates: *(i)* Random Regular Graphs (RRG), in which all the nodes have the same number of contacts ($k_i = \langle k\rangle, \forall i$), and *(ii)* Erdős-Rényi (ER) networks [31], for which the probability of finding a node with $k$ neighbors, $P(k)$, follows a Poisson distribution so that most of the nodes have a degree $k$ close to the average $\langle k\rangle$. Besides, we have analyzed networks having a scale-free (SF) pattern for the probability distribution, $P(k) \sim k^{-3}$, constructed by means of a configurational random model (SF-CONF) [32] and the Barabási-Albert model (SF-BA) [33].

*Macroscopic analysis.-* First we report the average entanglement entropy per node $\langle S_i\rangle = \sum_{i=1}^{N} S_i/N$, as a function of the coupling between oscillators. From Fig. 1 it is clear that entanglement increases with the coupling strength $c$. However, fixing the value of the coupling, the entropy per node of each network shows differences. In particular, as the heterogeneity of the degree distribution $P(k)$ of the network increases (from RRG to SF networks) the value of $\langle S_i\rangle$ decreases. Thus, the presence of different connectivity classes within the same network seems to be detrimental for the (average) entanglement within its components.

*Microscopic analysis.-* The above finding motivates the study of the entanglement entropy at the level of single nodes in networks having different connectivity classes such as ER and SF networks. To this aim, we collect the entanglement entropies of the $N_k$ nodes having con-



FIG. 2. (color online) In **(a)** we show the microscopic picture: a particular node and its boundary (first neighbors) with the rest of the system (blurred). The rest of the panels show the average entropy of nodes with degree $k$, $\langle S_k \rangle$ for the following topologies: **(b)** ER, **(c)** SF-CONF and **(d)** SF-BA networks. Each symbol refers to a value of the coupling strength $c$. In particular, we have that ●,■,◆,▲ correspond to $c = 0.2$, 0.4, 0.6 and 0.8 respectively. All the networks have the same average degree $\langle k \rangle = 4$. Solid lines represent the theoretical curves calculated using the mean-field formulation.

nectivity $k$ and define the average entanglement of the degree class $k$ as: $\langle S_k \rangle = \sum_{i|k_i=k} S_i/N_k$. One would expect that the larger the connectivity $k$ of a node the more correlated with the rest of the network it is, and thus the larger the value of $\langle S_k \rangle$ is.

The growth of $\langle S_k \rangle$ with $k$ would point out the existence of an area law [34] for complex networks. The area law states that the entanglement entropy of an inner region of a system scales with the size of the boundary connecting it with the rest of the system. For a regular lattice, the size of the boundary of an element is given by twice its dimensionality thus, in analogy, for a node in a complex network its boundary [see Fig. 2.**(a)**] is given by its connectivity, *i.e.*, $\langle S_k \rangle \sim k$. Taking this picture, one can consider a heterogeneous network as a collection of different dimensionalities within the same system.

The panels in Figs. 2 and 3 summarize our findings for the behavior of $\langle S_k \rangle$. In Fig, 2 we explore ER [panel **(b)**], SF-CONF [panel **(c)**] and SF-BA [panel **(d)**] networks for different values of the coupling $c = 0.2, 0.4, 0.6$ and 0.8. For ER networks we observe that the value of $\langle S_k \rangle$ increases with $k$ although we note that the growing trend seems to saturate for large values of $k$ pointing out that entanglement is bounded. On the other hand for SF networks, displaying a larger heterogeneity for the collection of degrees, the growing trend of $\langle S_k \rangle$ only holds for small to moderate connectivities $k$, then $\langle S_k \rangle$ reaches a maximum and starts to decrease. Eventually, those nodes with sufficiently large $k$ would drop its entanglement entropy. As a result, *hubs* are not the most entangled nodes, but there is an optimally-correlated class of nodes having moderate connectivity. The plots in Fig. 3 confirm the above results. In these cases, we have fixed $c = 0.6$ and changed the mean connectivity of the RRG [panel **(a)**], ER [panel **(b)**], SF-CONF [panel **(c)**] and SF-BA [panel **(d)**] networks [35].



FIG. 3. (color online) Average entropy of nodes with degree $k$, $\langle S_k \rangle$, for all the topologies under study: **(a)** RRG , **(b)** ER, **(c)** SF-CONF and **(d)** SF-BA networks. All the oscillators are coupled with the same strength $c = 0.6$. Each symbol represents a different value of the average degree $\langle k \rangle$ of the system. Solid lines represent the theoretical behavior calculated with the mean-field formulation.

*Mean-field formulation.-* In the following we develop a minimal model aimed at capturing the rise-and-fall behavior of $\langle S_k \rangle$ in SF networks. We propose a generalized mean field approximation that goes beyond the standard one in which the many-body Hamiltonian is reduced by replacing the interaction terms by their mean value, thus hindering the effect of the heterogeneity in the number of inputs (degree) that nodes receive.

The simplest framework for dealing with bipartite entanglement is shown in Fig. 2.**(a)**. We consider both the node of interest (say 0) and its first $k$ neighbors, whereas the influence of the rest of the network (blurred in the figure) is taken into account by renormalizing the frequency of the latter ones. In this approximation, the local Hamiltonian of a node of degree $k$ reads,

$$H^k_{\mathrm{MF}} = \frac{1}{2}\left\{ p_0^2 + x_0^2 + \sum_{j=1}^{k} \left[ p_j^2 + \nu_\kappa^2 x_j^2 + c(x_0 - x_j)^2 \right] \right\},$$

where the renormalized frequency of the $k$ neighbbors reads: $\nu_\kappa^2 = 1 + c^2\kappa$, where $\kappa$ is a fitting parameter (see below).

This model is exactly solvable (see Appendix B). Therefore, we can find the entropy of the central node $\langle S_k \rangle^{MF}$ analytically as a function of its connectivity $k$ and $\kappa$. In Figs. 2 and 3 we plot with solid lines the curves $\langle S_k \rangle^{MF}$ obtained after tuning the single parameter $\kappa$ for each of the curves. Interestingly, the values of $\kappa$



FIG. 4. (color online) Controlled entanglement via an external (blue) node coupled to an ER graph. In the plot we show (filled dots) the entanglement $\langle S_k \rangle$ of the nodes of the ER network as a function of their connectivity. In addition, we show (empty squares) the evolution of the entanglement of the external controller as a function of its connectivity, *i.e.* the number of links launched to the target network.

used for each curve only depends on the network topology. As reported in Appendix C, its value is independent of the value of $c$ but proportional to the average conectivity $\langle k \rangle$. From the figures it is clear that our mean-field approximation agrees fairly good with numerics.

In addition to the quantitative agreement, the analytical estimation $\langle S_k \rangle^{MF}$ allows to explain the rise-and-fall of entropy across degree classes. As shown in the Appendix C this phenomenon lies in the fact that hubs are *almost* eigenvectors of the Laplacian and thus normal modes of the Hamiltonian (uncoupled from the rest of the system and therefore not entangled). The progressive localization of the eigenvectors with $k$ balances the growth of the correlations associated to the increase of $k$. It is the competition between these two effects what explains the peak for $\langle S_k \rangle$ in SF networks at moderately, rather that maximally, coupled nodes.

*Controlling entanglement.-* The results presented above illustrate the particular properties that quantum networks introduce with respect to its classical counterparts. To round off our discussion, we want to connect the rise-and-fall of the entanglement entropy with the relevant issue of the control of networks [36] and span it to the quantum domain. Let us suppose that we have a quantum network with a rather regular topology, *e.g.* and ER graph. Our aim is to obtain the maximum entanglement with the network by adding an external node (our controller) as shown in Fig. 4. Two relevant questions arise: *(i)* how many connections should we launch to the system under control? and *(ii)* what are the target nodes to be connected with the controller?

In a random network, such as an ER graph, almost all the nodes are equivalent, thus we can assume that there are not preferred nodes to attach with, so that the number of optimal connections remains as the most relevant issue to address. One would be tempted to state that the more connections we add the more entanglement with the system the controller will obtain. However, in Fig. 4 we show that the entanglement obtained by the external controller increases up to a maximum at $k \simeq 15$, while a further increase of its connectivity leads to the decrease of its entanglement entropy. The experiment illustrates that the amount of information we can extract from a quantum network via an external controller is bounded and its maximum is reached by means of a moderate number of connections with the target system.

*Conclusions.-* The entropy of entanglement of nodes in networks of quantum oscillators reveals a non-trivial behavior. The decay of the entanglement for large connectivity nodes is seen as the fingerprint of the localization of some of the Laplacian eigenvectors around hubs which turns them into normal modes of the system. This effect balances the increase of the entanglement with the connectivity, analogously to an area law in regular lattices, thus causing the rise-and-fall of the entanglement entropy across connectivity classes. As a byproduct, the control of quantum networks by maximizing the entanglement with an external controller is limited and does not rely in maximizing the number of physical connections with the network under study.

In addition to their interest for the emerging field of quantum information networks, our results show an interesting connection with fundamental concepts of quantum gravity in complex spatial connectivities. In fact, the setup used here can be seen as the discretized version of real massive Klein-Gordon fields far from the usual Minkowsky or curved space-time situations, suggested from emergent gravity concepts as intermediate topologies in the transition from a highly connected (high energy) quantum geometric phase of the universe to the low energy, largely homogeneous, actual phase [37, 38]. Although the connection between complex space-time topologies and the field of network science has recently attracted attention [39], the above scenario paves the way for the study of quantum models on complex space-time topologies thus fostering the synergy between these apparently unrelated fields of physics.

## ACKNOWLEDGMENTS

We acknowledge Diego Blas for discussions and support from the Spanish DGICYT under projects, FIS2011-23526 and FIS2011-25167, and by the Aragón (Grupo FENOL) and Balearic Goverments. F.G. acknowledges the CSIC post-doctoral JAE program. J.G.G. is supported by MICINN through the Ramón y Cajal program.

### Appendix A: Entanglement entropy

Let us start by deriving Eq. (4) in the text. Given $\varrho$, the quantum state of the network, its associated Von

Neumann entropy is given by,

$$S = -\mathrm{Tr}(\varrho \ln \varrho)\ , \tag{A1}$$

where Tr accounts for the trace operation. The marginal entropy for the node $i$, $S_i$, is obtained by replacing $\varrho$ by the reduced density matrix,

$$\varrho_i = \mathrm{Tr}_{i^c}\,\varrho\ , \tag{A2}$$

in (A1) with $\mathrm{Tr}_{i^c}$ the partial trace. The latter means the trace over the complement of $i$ (the rest of the nodes).

In our work, the state of the network considered is the ground state of the Hamiltonian [Eq. (1) in the text]:

$$H_{\mathrm{network}} = \frac{1}{2}\Big(\mathbf{p}^{\mathrm{T}}\mathbf{p} + \mathbf{x}^{\mathrm{T}}(\mathbb{I} + 2cL)\mathbf{x}\Big)\ , \tag{A3}$$

here $\mathbb{I}$ is the $N \times N$ identity matrix, $c$ is the coupling strength between connected oscillators and $L$ the network laplacian. The operators $\mathbf{p}^{\mathrm{T}} = (p_1, p_2, ..., p_N)$ and $\mathbf{x}^{\mathrm{T}} = (x_1, , x_2, ..., x_N)$ are the momenta and positions of nodes respectively, satisfying the usual commutation relations: $[\mathbf{x}, \mathbf{p}^{\mathrm{T}}] = \mathrm{i}\hbar\mathbb{I}$. We are interested in analyzing the ground state of the system, which is a pure state, thus having $S = 0$. On top of that, this state is Gaussian (since the Hamiltonian is quadratic) so that the reduced density matrices $\varrho_i$ and $S_i$ can be computed by means of the covariance matrix:

$$\sigma = \begin{pmatrix} \langle x_i^2\rangle & \frac{1}{2}\langle x_i p_i + p_i x_i\rangle \\ \frac{1}{2}\langle x_i p_i + p_i x_i\rangle & \langle p_i^2\rangle \end{pmatrix}\ , \tag{A4}$$

where the averages are calculated via the reduced density matrix, $\varrho_i$, as: $\langle x_i^2\rangle = \mathrm{Tr}(x_i^2 \varrho_i)$.

It was Agarwal [30] who derived an explicit formula for the marginal entropies [Eq. (3) in the text]:

$$S_i = S_{i^c} = \Big(\mu_i + \frac{1}{2}\Big)\log\Big(\mu_i + \frac{1}{2}\Big) - \Big(\mu_i - \frac{1}{2}\Big)\log\Big(\mu_i - \frac{1}{2}\Big)\ , \tag{A5}$$

with $\mu_i = \sqrt{\langle x_i^2\rangle\langle p_i^2\rangle}$.

We are able to can find these quadratures by working with normal modes, *i.e.*, those diagonalizing the potential energy matrix $V = \mathbb{I} + 2cL$:

$$\vec{x} = S\vec{Q} \text{ so that } S^T V S = V_d, \tag{A6}$$

whose quadratures are those of a set of uncoupled oscillators at their individual ground state:

$$\langle Q_i\rangle = \langle P_i\rangle = 0 \tag{A7}$$

$$\langle Q_i Q_j\rangle = \delta_{ij}\frac{\hbar}{2\Omega_j} \tag{A8}$$

$$\langle P_i P_j\rangle = \delta_{ij}\frac{\hbar\Omega_j}{2} \tag{A9}$$

$$\frac{1}{2}\langle Q_i P_j + Q_j P_i\rangle = 0, \tag{A10}$$

with $\Omega_i = \sqrt{1 + 2c\lambda_i}$ the eigenfrequencies, and $\lambda_i$ the eigenvalues of the laplacian matrix $L$. The latter relation is obtained from simple inspection of the eigenvalue equation $V\vec{u} = \Omega^2\vec{u} = (\mathbb{I} + 2cL)\vec{u}$, so that $L\vec{u} = \frac{\Omega^2 - 1}{2c}\vec{u} \equiv \lambda\vec{u}$. That is, $\vec{u}$ is an eigenvector of both $V$ and $L$, with $\Omega = \sqrt{1 + 2c\lambda}$. Then we obtain:

$$\langle x_i\rangle = \sum_j S_{ij}\langle Q_j\rangle = 0 \tag{A11}$$

$$\langle p_i\rangle = \sum_j S_{ij}\langle P_j\rangle = 0 \tag{A12}$$

$$\langle p_i x_j\rangle = \sum_{k,l} S_{ik} S_{jl}\langle P_k Q_l\rangle = 0 \tag{A13}$$

$$\langle x_i^2\rangle = \sum_j \big(S_{ij}\big)^2\langle Q_j^2\rangle = \sum_j \big(S_{ij}\big)^2\frac{\hbar}{2\Omega_j} \tag{A14}$$

$$\langle p_i^2\rangle = \sum_j \big(S_{ij}\big)^2\langle P_j^2\rangle = \sum_j \big(S_{ij}\big)^2\frac{\hbar\Omega_j}{2}. \tag{A15}$$

Finally, we arrive to the expression for the quadratures:

$$\begin{aligned} \mu_i^2 &= \langle x_i^2\rangle\langle p_i^2\rangle - \frac{1}{2}\langle x_i p_i + p_i x_i\rangle \\ &= \langle x_i^2\rangle\langle p_i^2\rangle = \frac{1}{4}\sum_j \big(S_{ij}\big)^2\big(S_{ij'}\big)^2\frac{\Omega_j}{\Omega_{j'}}, \end{aligned} \tag{A16}$$

as stated in the main text.

## Appendix B: Entanglement mean field approximation

In this section we sketch the solution for the mean field approximation presented in the main text. The mean-field Hamiltonian for a node surrounded by $k$ neighbors [Cf. Fig. 1.**a** in the article]can be rewritten in matrix form like as:

$$H = \frac{1}{2}\Big(\mathbf{p}^t\,\mathbb{I}\,\mathbf{p} + \mathbf{x}^t\,\hat{V}\,\mathbf{x}\Big) \tag{B1}$$

whith the (k+1)-tuples:

$$\mathbf{p} = \begin{pmatrix} p_0 \\ p_1 \\ \vdots \\ p_k \end{pmatrix}\ , \qquad \mathbf{x} = \begin{pmatrix} x_0 \\ x_1 \\ \vdots \\ x_k \end{pmatrix}\ , \tag{B2}$$

note that we have named the 0-node the central one. The then potential reads:

$$\hat{V} = \begin{pmatrix} 1 + 2c\,k & -2c & -2c & \cdots & -2c \\ -2c & 1 + 2c\,\kappa & 0 & \cdots & 0 \\ -2c & 0 & 1 + 2c\,\kappa & \cdots & 0 \\ \vdots & \vdots & & \ddots & \vdots \\ -2c & \cdots & \cdots & 0 & 1 + 2c\,\kappa \end{pmatrix} \tag{B3}$$

The equilibrium properties of (B1), in particular the Von Neumann entropy, is characterized by the eigenvalues and eigenvectors of $\hat{V}$, as explained in the previous section. It turns out that the spectrum of $\hat{V}$ given by (B3) can be analytically computed:

*i)* The $(k+1)\times(k+1)$ matrix $\hat{V}$ in (B3) has $(k+1)-2$ eigenvectors of the form:

$$|\lambda_j\rangle = \begin{pmatrix} 0 \\ \vdots \\ 1_i \\ \vdots \\ -1_j \\ \vdots \end{pmatrix} \tag{B4}$$

with degenerated eigenvalues

$$\lambda_j = 1 + 2c\kappa \tag{B5}$$

as can be easily checked.

*ii)* The other two eigenvectors are of the form,

$$|\lambda\rangle = \frac{1}{\sqrt{z^2+k}} \begin{pmatrix} z \\ 1 \\ \vdots \\ 1 \end{pmatrix} \tag{B6}$$

where the eigenvalues and eigenvectors are found from the equations,

$$(1 + 2ck^2) - kc^2 = \lambda z \ , \tag{B7}$$

$$-c^2 z + 1 + 2c\kappa = \lambda \ , \tag{B8}$$

with eigenvalues

$$\lambda_\pm = \frac{1}{2}\left\{2c\left[\kappa + \sqrt{\kappa^2 - 2\kappa k + k(k+4)} + k\right] + 2\right\} \tag{B9}$$

and

$$z_\pm = \frac{1}{2}\left[\kappa - k \mp \sqrt{\kappa^2 - 2\kappa k + k(k+4)}\right] \tag{B10}$$

The latter are the only ones entering in the formula for the marginal entropy of the node 0, see (A5) and Eq. (3) in maint text. Thus, the quadratures can be written as:

$$\langle x_0^2\rangle = \sum_{i=\pm} \frac{z_i^2}{z_i^2+k}\frac{1}{2\sqrt{\lambda_i}} \ , \tag{B11}$$

$$\langle p_0^2\rangle = \sum_{i=\pm} \frac{z_i^2}{z_i^2+k}\frac{\sqrt{\lambda_i}}{2} \ , \tag{B12}$$

from which the entropy is obtained.

We finally note that $z_+ \to -k$ for large enough $k$. Therefore in this limit the corresponding eigenvector approaches to $(1, 0, ..., 0)$ with frequency $\omega = \sqrt{\lambda_+} \to \sqrt{1+2ck}$. Therefore, in the limit of large connectivity the node is a normal mode and its corresponding marginal entropy approaches to zero.

## Appendix C: Extimation of fitting parameter $\kappa$ values

Finally, we focus on the values of the fitting parameter $\kappa$ used to compute the mean-field solution shown in Figs. 2 and 3 of the main text. The values of $\kappa$ for the different topologies under study are shown in Tab. I. As observed, in the case of Erdős Rényi networks with average degree $\langle k\rangle = 2$ results are missing. This is due to the fact that for average degrees lesser than 2.5 the resulting ER networks are not made by a unique connected component. As already stated in the main text, we observe that $\kappa$ only depends on the average degree and on the considered underlying topology. Also, as a further validation of our mean field hypotesis, with the only exception of Scale-Free networks made through the configurational model with $\langle k\rangle = 4$, we notice that $\kappa$ always assumes values slightly below the average degree.

| $\langle k\rangle$ \ $c$ | **RRG** | **ER** | **SF-BA** | **SF-CONF** |
|---|---|---|---|---|
| 2 \0.2 | – | 2.6 | 1.8 | 2.5 |
| 2 \0.4 | – | 2.6 | 1.8 | 2.5 |
| 2 \0.6 | – | 2.6 | 1.8 | 2.5 |
| 2 \0.8 | – | 2.6 | 1.8 | 2.5 |
| 4 \0.2 | 3.4 | 3.8 | 3.5 | 4.2 |
| 4 \0.4 | 3.4 | 3.8 | 3.5 | 4.2 |
| 4 \0.6 | 3.4 | 3.8 | 3.5 | 4.2 |
| 4 \0.8 | 3.4 | 3.8 | 3.5 | 4.2 |
| 6 \0.2 | 5.2 | 5.5 | 5.2 | 5.8 |
| 6 \0.4 | 5.2 | 5.5 | 5.2 | 5.8 |
| 6 \0.6 | 5.2 | 5.5 | 5.2 | 5.8 |
| 6 \0.8 | 5.2 | 5.5 | 5.2 | 5.8 |

TABLE I. Fitting parameter $\kappa$ for all the different network topologies with respect to each pair of parameters average degree $\langle k\rangle$ and coupling $c$.

[1] A.L. Barabási, Nature Phys. **8**, 14 (2012).

[2] R. Albert and A.-L. Barabási, Rev. Mod. Phys. **74**, 47 (2002).

[3] M.E.J. Newman, SIAM Rev. **45**, 167 (2003).
[4] M.E.J. Newman, *Networks: An Introduction*, Oxford University Press, USA (2010).
[5] S. Boccaletti, V. Latora, Y. Moreno, M. Chavez and D.-U. Hwang, Phys. Rep. **424**, 175 (2006).
[6] A. Vespignani, Nature Phys. **8**, 32 (2012).
[7] G. Bianconi, EPL **81**, 28005 (2008).
[8] G. Bianconi, Phys. Rev. E **79**, 036114 (2009).
[9] K. Anand and G. Bianconi, Phys. Rev. E **80** 045102 (2009). topologies
[10] S. Johnson, J.J. Torres, J. Marro and M.A. Muñoz, Phys. Rev. Lett. **104**, 108702 (2010).
[11] G. Bianconi, P. Pin and M. Marsili, Proc. Nat. Acad. Sci. (USA) **106**, 11433 (2009).
[12] J. Gómez-Gardeñes and V. Latora, Phys. Rev. E **78**, 065102 (2008).
[13] Z. Burda, J. Duda, J.M. Luck and B. Waclaw, Phys. Rev. Lett. **102**, 160602 (2009).
[14] R. Sinatra, J. Gómez-Gardeñes, R. Lambiotte, V. Nicosia and V. Latora, Phys. Rev. E **83**, 030103 (2011).
[15] M. Rosvall and C.T. Bergstrom, Proc. Nat. Acad. Sci. (USA) **104**, 7327 (2007).
[16] M. Rosvall and C.T. Bergstrom, Proc. Nat. Acad. Sci. (USA) **105**, 1118 (2008).
[17] A. Acín, J.I. Cirac and M. Lewenstein, Nature Phys. **3**, 256 (2007).
[18] M. Cuquet and J. Calsamiglia, Phys. Rev. Lett. **103**, 240503 (2009).
[19] M. Cuquet and J. Calsamiglia, Phys. Rev. A **83**, 032319 (2011).
[20] L. Wu and S.Q. Zhu, Phys. Rev. A **84**, 052304 (2011).
[21] S. Perseguers, M. Lewenstein, A. Acín and J.I. Cirac, Nature Phys. **6**, 539 (2010).
[22] O. Muelken and A. Blumen, Phys. Rep. **502**, 37 (2011).
[23] G.D. Paparo and M.A. Martín-Delgado, Sci. Rep. **2**, 444 (2012).
[24] S. Garnerone, P. Zanardi and D.A. Lidar, Phys. Rev. Lett. **108** 230506 (2012).
[25] S Garnerone, Phys. Rev. A **86**, 032342 (2012)
[26] E. Sánchez-Burillo, J. Duch, J. Gómez-Gardeñes and D. Zueco, Sci. Rep. **2** 605 (2012).
[27] S.L. Braunstein, S. Ghosh, T. Mansour, S. Severini and R.C. Wilson, Phys. Rev. A **73**, 012320 (2006).
[28] K. Anand, G. Bianconi and S. Severini, Phys. Rev. E **83**, 036109 (2011).
[29] P. Van Mieghem, *Graph Spectra for Complex Networks*, Cambridge University Press, UK (2010).
[30] G. Agarwal, Phys. Rev. A **3**, 828 (1971).
[31] P. Erdős and A. Rényi, Publ. Math. Inst. Hung. Acad. Sci. **5**, 17 (1960).
[32] M. Molloy and B. Reed, Random Struct. Algorithms **6**, 161 (1995).
[33] A.L. Barbási and R. Albert, Science **286**, 509 (1999).
[34] J. Eisert, M. Cramer and M. B. Plenio, Rev. Mod. Phys. **82**, 277 (2010).
[35] Note that for the panel 2.**a** corresponding to RRG, each of the points corresponds to an ensemble of graphs in which all the nodes have the same connectivity $k$. The decreasing trend points out that for this (single-scale) class of graphs entanglement decreases with connectivity for the values of $k$ considered.
[36] Y.-Y. Liu, J.-J. Slotine and A.-L. Barabási, Nature **473**, 167 (2011).
[37] C. Prescod-Weinstein and L. Smolin, Phys. Rev. D **80**, 063505 (2009).
[38] F. Markopoulou and L. Smolin, Class. Quantum Grav. **24**, 3813 (2007).
[39] D. Krioukov, M. Kitsak, R.S. Sinkovits, D. Rideout, D. Meyer and M. Boguñá, *Network Cosmology* arXiv:1203.2109.